

University of Dundee

## Skin cancer burden in lung transplant recipients

Proby, C. M.; Harwood, C. A.

*Published in:*  
British Journal of Dermatology

*DOI:*  
[10.1111/bjd.18903](https://doi.org/10.1111/bjd.18903)

*Publication date:*  
2020

*Licence:*  
CC BY-NC

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

*Citation for published version (APA):*  
Proby, C. M., & Harwood, C. A. (2020). Skin cancer burden in lung transplant recipients: we need to do better! *British Journal of Dermatology*, 183(3), 416-417. <https://doi.org/10.1111/bjd.18903>

### General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Skin cancer burden in lung transplant recipients: we need to do better!

DOI: 10.1111/bjd.18903

Linked Article: Way et al. *Br J Dermatol* 2020; DOI: 10.1111/bjd.18812

Solid organ transplant recipients (SOTRs) are at significantly increased risk for multiple skin cancers.<sup>1</sup> Skin cancer multiplicity matters: immunosuppressed patients with multiple squamous cell carcinomas (SCCs) have poorer outcomes.<sup>2</sup> Routine skin cancer surveillance and preventative strategies have been widely adopted in many centres, but current practice may fall short for lung transplant recipients (LTRs).

In this issue of the *BJD*, Way and colleagues<sup>3</sup> quantify the burden of new skin cancers developing annually in LTRs, and the numbers are staggering. This study comes from Queensland, Australia, arguably the 'skin cancer capital' of the world, but also a region with sophisticated strategies to combat skin cancer in the general population, and with high levels of education among patients and general practitioners.

High incidence rates for keratinocyte cancers (KCs) in SOTRs have been extensively reported, usually based on first primary KC. Here the authors assess total burden of KCs in a prospective, population-based study. Eligibility was restricted to patients stably transplanted for at least 1 year in whom systemic retinoids or topical therapies had not been initiated in the previous 6 months and 125 LTRs in Queensland completed the study. By choosing a population-based study and by systematically cross-checking both public and private pathology databases for confirmation of interval skin cancers, these authors established age-standardized incidence rates of 447 per 1000 patient-years for SCC and 281 per 1000 person-years for basal cell carcinoma (BCC). This is more than double the age-standardized incidence rates for LTR SCC when based on first primary SCC alone (201 per 1000 person-years) and is 77 times higher than the estimated SCC incidence rate in the general Queensland population.

Why do LTRs have such a high skin cancer risk? As in cardiac transplantation, LTRs need greater levels of immunosuppression than renal or liver transplant recipients. This intense immunosuppression interacts with previous and ongoing solar ultraviolet (UV) exposure, creating a perfect storm for skin carcinogenesis. LTRs have historically received azathioprine as part of their maintenance immunosuppressive regimen, although mycophenolate mofetil is now preferentially used. Many LTRs also receive voriconazole for treatment or prevention of fungal lung infections. Like azathioprine,<sup>4,5</sup> voriconazole is photosensitizing to UVA<sup>6</sup> and contributes to carcinogenesis both directly and indirectly through mechanisms that include generation of reactive oxygen species.

When examining risk factors associated with skin cancer multiplicity, Way and colleagues found that ever use of voriconazole increased risk of SCC (not BCC) twofold, and treatment with voriconazole for 4 or more months increased SCC risk 4-5-fold. Thus, regimens that include prophylactic voriconazole will compound the already very high risk for cutaneous SCC in LTRs.

As dermatologists, we are ideally placed to provide dedicated OTR skin clinics for long-term skin surveillance, rapid access for urgent problems and skin cancer risk reduction management. Such opportunities are often limited during transplant clinic consultations, in which many other clinical problems compete for the transplant team's time and resources. Nonetheless, we should be aiming to develop skin cancer prevention programmes for routine delivery within transplantation clinics. Simple protocols, ideally using multimodal methods, could be created to support best practice in sun protection and skin cancer education. Such approaches are a focus of recent research efforts and may be a starting point for addressing the huge and growing burden of OTR skin cancer so dramatically highlighted in this study.

C.M. Proby<sup>1</sup> and C.A. Harwood<sup>2</sup>

<sup>1</sup>Division of Molecular and Clinical Medicine, Ninewells Hospital and Medical School, University of Dundee, Dundee, DD1 9SY, UK and <sup>2</sup>Centre for Cell Biology and Cutaneous Research, Blizard Institute, Queen Mary University of London, London, E1 2AT, UK  
Email: c.proby@dundee.ac.uk

Conflicts of interest: none to declare.

## References

- Garrett GL, Blanc PD, Boscardin J et al. Incidence of and risk factors for skin cancer in organ transplant recipients in the United States. *JAMA Dermatol* 2017; **153**:296–303.
- Gonzalez JL, Reddy ND, Cunningham K et al. Multiple cutaneous squamous cell carcinoma in immunosuppressed vs immunocompetent patients. *JAMA Dermatol* 2019; **155**:625–7.
- Way M, Marquart L, Chambers DC et al. Skin cancer multiplicity in lung transplant recipients: a prospective, population-based study. *Br J Dermatol* 2020; <https://doi.org/10.1111/bjd.18812>. [Epub ahead of print]
- Leigh IM, Proby CM, Inman GJ, Harwood CA. Azathioprine: friend or foe? *Br J Dermatol* 2019; **180**:961–3.
- Jiyad Z, Olsen CM, Burke MT et al. Azathioprine and risk of skin cancer in organ transplant recipients: systematic review and meta-analysis. *Am J Transplant* 2016; **16**:3490–503.
- Ona K, Oh DH. Voriconazole N-oxide and its ultraviolet B photoproduct sensitise keratinocytes to ultraviolet A. *Br J Dermatol* 2015; **173**:751–9.